

BIABA

Composition: abamectin 1,8% EC p/v

ACARIZIDE INSECTICIDE

Formulation with insecticide and acaricide characteristics, to be used in foliar pulverization to control plagues in several cultivations.

Emulsionable-concentrated composition used to control psylla on pear trees, attacker of citrus trees and other undermine caterpillar, *Liriomyza* (underwater) and other Dipterous undermine larva, as well as pear trees erynosis and tetra-niquides mobile shapes (red and yellow spiders) and tarsus-nemides (white spiders) in cultivations, vegetable plantations and hotbed, all of them mentioned in every specific assumption.

Application doses:

	Doses	Observations
Cottoner	0,05-0,10 %	Mites
Celery, cucurbitaceae, tomato, pepper	0,05-0,10 %	Mites and Liryomize
Ornamentals, herbaceous, strawberry bed	0,05-0,10 %	Mites and Liryomize
Citrus trees	0,05-0,10 %	Phylcnistis
Pear tree	0,05-0,10 %	Mites, erinosys and psila
Mulberry, raspberry	0,05-0,10 %	Red spider

In every case, recommended doses must be applied in normal pulverization way, no exceeding 1'5 l/ha and application. In woody cultivations, some technicians recommend to add 200-500 cc oil in summertime/Hl. of broth and starting treatments when new individuals turn up. In order to make more difficult the showing up of resistances in mites, it is recommended no treating more than 3 times a year and alternate with different application acaricides.

Subscribed on R.O.P.M.F. N° 24.070/16



AGROSANITARIO S.L.

Apdo. Correos 2019. 30509. Molina de Segura (Murcia).
Tel y fax. 0034 968 619 515. www.agrosanitario.com